ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:

6-benzylaminopurine (aytokinin)/ 6-bensietaminopurien (sitokinien)._____ 21 g/c

Registration holder / Registrasiehouer: UNIVERSAL CROP PROTECTION (PTY) LTD Co. Reg. No. 1983/008184/07 Mpv. Reg. Nr. RO, Box / Posbus 801 KEMPTON PARK, 1620 T(011) 396 2233 Website /

Webblad: www.villacrop.co.za

Manufactured by / Veryaardig deur,

grochemicals Limited. Hill End House. Whittington, Worcester, United Kingdom

Vir valledige besonderhede, sien ingeslote pamilet.





Reg. No. L 7891 Act/No. 36 of/van 1947

A water-soluble concentrate plant growth regulator for fruit thinning on apples. 'n Wateroplosbare konsentraat plantgroeireguleerder vir uitdun van vrugte op appels.

Net contents / Netto inhoud: 56

> UN number: Not Regulated

Date formulated: Eormuleringsdatum: Batch No / Lotnr:

Distributed by / Versprei deur:



Tel: (011) 396 2233

EXILISG is the registered trademark of FINE Agrochemicals Ltd., UK. EXILES® is die geregistreerde handelsmerk van FINE Agrochemicals Ltd., VK.





CAUTION VERSIGTIG





. Harmful if swallowed.

WARNINGS

Store in a cool place, away from food, feeds, seed, fertilizer and other agricultural remedies.

Can cause eve irritation.

Keep out of reach of children, uninformed persons and animals. Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

registration. Consult the supplier in event of any uncertainty.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the

label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the

PRECAUTIONS

- Do not inhale the spray mist.
- Avoid contact with skin, eyes and clothing.
- Wash with soap and water after use.
- In case of contact with eyes, flush with plenty of water and get medical attention.
- In case of contact with skin, wash immediately with soap and plenty of water and remove contaminated clothes. Do not eat, drink or smoke whilst mixing or applying the product or before washing hands and face or change of
- Prevent drift onto other crops, grazing, rivers, dams or areas not under treatment.
- Clean the applicator after use and dispose of wash water where it will not contaminate food, grazing, rivers or dams.
- TRIPLE RINSE empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinsing the container three times with a volume of water equal to a minimum of 10 % of that of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the container by perforation and flattening and dispose of it in a safe way.
- Never re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.
- DIRECTIONS FOR USE; Use only as directed.

Compatibility:

- Do not apply EXILIS® in combination with pesticides and other growth regulators.
- It is advisable to add a buffer e.g. COMMODOBUFF (L. 5390) to alkaline spray water to reduce the pH below 7.

Application Instructions:

- EXILIS® is a plant growth regulator for use on apples as a post-bloom thinning agent.
- EXILIS® must be absorbed by the plant to be effective.
- Optimum results will be achieved on vigorous, healthy trees when sprayed under good absorption conditions.
- High relative humidity and slow drying conditions favour maximum absorption. Do not apply EXILIS® to low vigor trees, or trees under stress from drought, winter injury etc.
- Apply as a uniform fine mist spray so that 85 % of the spray volume is directed into the upper two thirds of the trees
- under calm weather conditions. Use a property calibrated sprayer that ensures uniform and complete coverage of all foliage and blossoms. Spray volume should be adjusted to tree size and density. Spray volumes around 1000 litres per hectare should
- Do not apply EXILIS® when the air temperature is lower than 10 °C or greater than 32 °C. Rapid drying conditions should be avoided. Applications in the mornings or evenings may be preferable.
- Temperatures of over 18°C for the two days following application are considered important for good thinning. High temperatures (+28 °C over this period) may result in excessive thinning.

- Deay EXILISE application after rain until lower parts and leaves are free of surface droplets. Hainfall within 6 hours of application may reduce the expected response of EXILIS®.
- Thinning responses are normally to be observed within 10 to 14 days of application.

Mixing Instructions:

- Add the required amount of EXIUS® to a clean spray tank half filled with water.
- Agitate while filling the tank with water to the required level. EXILIS® spray solutions should be used within 24 hours of mixing.
- When a buffered wetting agent is used, it should be added to the tank prior to the addition of EXILIS®.

APPLICATION DOSAGE RATES:

Crop	Growth Regulating Activity	Dosage Rate mt / 100 t water	Remarks
Apples (Golden Delicious, Royal Gala, Pink Lady, Granny Smith)	Fruit Thinning	(in a minimum	Apply EXILIS® when the king fruitlets are between 10 to 12 mm, but not later than 30 days after full bloom.

EXILIS® is the registered trademark of/is die geregistreerde handelsmerk van FINE Agrochemicals Ltd., UK.

COMMODOBUFF is a registered product of / is 'n geregistreerde produk van Villa Crop Protection (Pty) Ltd.