

Crop / Use	Dosage Rate	Remarks
<b>Potatoes</b> Uncut tubers To improve sprouting of potato tubers.	Winter planting areas only. Variable results occur in the summer rainfall regions. 5 g /200 l water (2,5 g /100 l water)	Dip tubers for a period of 10 minutes in FALGRO® 20 SP solution. Leave in a cool dry place and allow tubers to dry. Plant as soon as first eye movement is observed. Change to a fresh solution after dipping sufficient tubers for planting one hectare. Do not dip cut tubers in this solution, as the results will be unsatisfactory.
Cut tubers To improve sprouting of potato tubers.	5 g /1000 l water	Dip tubers for a period of 10 minutes in solution after cutting. Leave in a cool dry place and allow tubers to dry. Plant as soon as first eye movement is observed. Change to a fresh solution after dipping sufficient tubers for planting one hectare. Do not dip uncut tubers in this solution, as results will be unsatisfactory.
<b>Forelle Pears</b> To improve fruit set.	5 g /100 l water	Apply as a full cover spray at 30 to 40 % flowering. Follow up with normal hand thinning 14 days after application.
<b>Citrus</b>		<ul style="list-style-type: none"> <li>Apply as a full cover spray. Do not apply to orchards where black spot has not been well controlled or where there is a high incidence of wind scar and Thrip markings. Do not apply to trees suffering from root disease or generally neglected trees. Do not allow trees to go into severe wilt.</li> <li>If a summer oil has been sprayed on citrus, a minimum of 14 days must elapse before FALGRO® 20 SP may be applied. An oil may be applied 5 to 10 days after a FALGRO® 20 SP application.</li> <li>Do not apply 2,4-D Ester to citrus destined for export before consulting the export guidelines.</li> </ul>
<b>Clementines</b> To increase fruit set.	1,25 g to 5 g /100 l water plus non-ionic wetting agent	Apply as a full cover spray from 65 to 75 % petal drop to 4 weeks after 100 % petal drop. With later applications applied, less influence on the fruit set reaction can be expected. The history of an orchard will influence FALGRO® 20 SP applications. Decide on the yield/fruit size balance before application. An excessive quantity of fruit will result in small fruit and induce alternate bearing. Get technical advice before applications are made.
<b>Grapefruit</b> Extension of on-tree storage period (delay of harvesting), larger and firmer fruit.	5 g /100 l water plus 2 ml 2,4-D Ester plus non-ionic wetting agent	Apply as a full cover spray at colourbreak/transparency (colour T5) or up to 3 weeks after colour break. Do not apply before colour break. Apply only to healthy trees where black spot and thrips are under control and no excessive wind scar occurs. The average fruit size must not be too large. The average diameter should exceed 89 mm at the time of spraying. Do not over-irrigate grapefruit trees intended for hanging.
<b>Lemons</b> To delay maturity (harvesting), thus to protect the crop during the decay prone period, fewer pickings and for larger fruits.	5 g /100 l water plus 2 ml 2,4-D Ester plus non-ionic wetting agent	Apply as a full cover spray 5 to 8 weeks before normal picking starts. Apply only to trees free of black spot or other blemishes.

Crop / Use	Dosage Rate	Remarks
<b>Citrus (continued)</b> Navels To reduce rind creasing in Navels oranges on Trifoliate or Trifoliate hybrid rootstock.	10 g /100 l water plus non-ionic wetting agent	Apply as a full cover spray in January to February but not later than the last week of February. Apply only to healthy trees where Thrip blemish and wind scars do not exceed 20 % cut on the total crop.
To reduce rind creasing in Navels Oranges on Rough lemon rootstock.	5 g /100 l water plus non-ionic wetting agent OR 5 g /100 l water plus 2 ml 2,4-D Ester plus non-ionic wetting agent	Apply as a full cover spray in January to February but not later than the last week of February. Apply only to healthy trees where Thrip blemish and wind scars do not exceed 20 % cut on the total crop. OR Apply as a full cover spray in May after full yellow colour has developed (minimum colour T3). Follow this treatment up with a 1 % urea (low biuret) at budbreak to avoid reduced flowering in the new season. Apply only to healthy trees where thrip blemish and wind scars do not exceed 20 % cut on the total crop.
<b>Valencia Oranges</b> To reduce rind creasing in Valencia oranges on all rootstocks	10 g /100 l water plus non-ionic wetting agent OR 5 g /100 l water plus non-ionic wetting agent OR 2,5 g /100 l water plus non-ionic wetting agent	Apply as a full cover spray in January to February but not later than the last week of February. Apply only to healthy trees where thrip blemish and wind scars do not exceed 20 % cut on the total crop. OR Apply as a full cover spray after the fruit has developed a full yellow colour (minimum colour T3). Do not apply before middle June. Apply only to healthy trees where thrip blemish and wind scars do not exceed 20 % cut on the total crop. OR Apply as a full cover spray when the fruit has developed a light greenish colour (T5 to T4, minimum colour T5). Do not apply before middle June. Apply only to healthy trees where Thrip blemish and wind scars do not exceed 20 % cut on the total crop.

FALGRO® is the registered trademark of / is die geregistreerde handelsmerk van Fine Agrochemicals Limited

ACTIVE INGREDIENT /  
AKTIEWE BESTANDDEEL:  
gibberellic acid  
.....200 g/kg  
gibberelinsuur

Manufactured by / Vervaardig deur:

fine

FINE AGROCHEMICALS LIMITED  
Hill End House, Whittington  
Worcester WR5 2RQ UK

E: enquire@fineagrochemicals.com  
www.fine-agrochemicals.com

FALGRO® is a registered trademark  
of / is die geregistreerde  
handelsmerk van:  
Fine Agrochemicals Limited.

UN number: Not Regulated

Vir volledige besonderhede,  
sien ingesluitte pamflet.  
For full particulars,  
see enclosed leaflet.



CAUTION  
VERSIGTIG



falgro® 20 SP

Reg. No. L 6394 Act/Wet No. 36 of 1947

A water-soluble powder plant growth hormone for the  
promotion of certain physiological and morphological  
effects as indicated on specific crops.

Packed in water-soluble sachets.  
'n Wateroplosbare poeier plant-groei-hormoon vir die  
beoordering van sekere fisiologiese en morfologiese  
effekte op spesifieke gewasse.  
Verpak in wateroplosbare sakke.

Net mass:  
Netto massa:  
(12 x 12,5 g)  
water-oplosbare sachette  
wateroplosbare sakke

UCP

Registration holder /  
Registrasiehouer  
UNIVERSAL CROP  
PROTECTION (PTY) LTD  
Co. Reg. No. 1903/06194/07  
Mpy. Reg. No.  
P.O. Box / Posbus 801,  
KEMPTON PARK, 1620  
Tel: (011) 306 2233  
Wobella / Wobbelad:  
www.villacrop.co.za

Distributed by/  
Versprei deur:  
VILLA CROP  
PROTECTION (PTY) LTD

VILLA  
CROP PROTECTION

Batch number:  
Lotnr:

Date of formulation:  
Formuleringsdatum:

FAL/W58/SA/16



**FALGRO® 20 SP**  
Reg. No.: L 6394 Act/Wet No. 36/1947

**ACTIVE INGREDIENT/ AKTIEWE BESTANDDEEL:**  
gibberellic acid / gibberelinsuur ..... 200 g/kg

Registration holder / Registrasiehouer:  
**UNIVERSAL CROP PROTECTION (PTY) LTD**  
Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.  
P.O. Box / Posbus 801, KEMPTON PARK, 1620, TEL: (011) 396 2233

**CAUTION / VERSIGTIG**

**WARNINGS**

- Handle the product with care,
- May cause mild skin irritation and slight eye irritation.
- Harmful when swallowed.
- Store in a cool, dry, well-ventilated place.
- Store away from food, feeds, seed, fertilizer and other agricultural remedies.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

**PRECAUTIONS**

- Do not inhale spray mist.
- Avoid contact with skin and eyes.
- Wash with soap and water after use and after accidental skin contact.
- Do not eat, drink or smoke whilst mixing or applying the product or before washing hands and face and change of clothing.
- Do not allow drift of spray onto other crops, grazing, rivers, dams or areas not under treatment.
- Packaging and containers must be completely emptied before being disposed of. Shake out thoroughly into the mixing tank and destroy the empty packaging in a safe way.
- Never re-use the empty packaging and containers for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.
- Clean the applicator and dipping bath after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams.

**USE RESTRICTIONS**

- The application of Gibberellic acid to certain citrus cultivars, such as Turkey Valencia and Bihahinna Naval, may cause phytotoxicity.
- Do not apply 2,4-D Ester in citrus destined for export before consulting the export guidelines.
- Do not use FALGRO® 20 SP on Sultanas, which are to be dried.
- Do not apply to vines, which are suffering from drought.

**DIRECTIONS FOR USE: Use only as directed.**

- Gibberellic acid performs most effectively when the spray/dip solution pH is 5,5 to 6,5. Higher pH values may impair product performance, whilst lower pH values may affect the crop negatively. It is recommended that the spray/dip solution be buffered, if the pH is outside 5,5 to 6,5.
- Environmental factors influence the efficacy of Gibberellic acid
- Water stress and heat waves can have negative influences.
- It is recommended that soil water conditions should be at field capacity during application.
- Applications should be early in the morning or late in the afternoon. Do not apply during the heat of the day.

**COMPATIBILITY:**

- Compatible with most normal spray materials, except for mineral oils and oil-in-water mixtures like Bordeaux mixture and Lime Sulphur.
- Do not use of borinated water.
- Do not apply with other agricultural chemicals not indicated on this label.
- Citrus: If a summer oil has been sprayed on citrus, a minimum of 14 days must elapse before FALGRO® 20 SP may be applied. An oil may be applied 5 to 10 days after a FALGRO® 20 SP application.

**MIXING INSTRUCTIONS:**

- Dissolve the required quantity of sachets in a small quantity of water.
- Ensure sachets are completely dissolved before adding the solution to the spray tank or dipping bath.
- Fill the spray tank/dipping bath with water to the required level, while maintaining agitation to ensure thorough mixing.
- Ensure the pH of the spray/dip solution is between 5,5 to 6,5. If not, add the suitable buffering agent to the solution before using.
- Maintain agitation during application.
- **IMPORTANT NOTE:** FALGRO® 20 SP is not very stable in water and spray mixtures should definitely not be left overnight.

**REFERENCE TO QUANTITY OF SACHETS PER DOSAGE RATE RECOMMENDATION:**

Quantity of FALGRO® 20 SP soluble sachets (12,5 g) required for different applications:

Dosage rate as indicated for specific uses on this label	Parts per million (ppm)	Dosage rate for various volumes of water	Quantity of soluble 12,5 g sachets required
1,25 g per 100 l water	2,5 ppm	12,5 g per 1000 l water	1 sachet per 1000 l water
2,5 g per 100 l water	5 ppm	12,5 g per 500 l water 25 g per 1000 l water	1 sachet per 500 l water 2 sachets per 1000 l water
5 g per 200 l water	5 ppm	25 g per 1000 l water	2 sachets per 1000 l water
5 g per 100 l water	10 ppm	25 g per 500 l water 50 g per 1000 l water	2 sachets per 500 l water 4 sachets per 1000 l water
10 g per 100 l water	20 ppm	50 g per 500 l water 100 g per 1000 l water	4 sachets per 500 l water 8 sachets per 1000 l water
15 g per 100 l water	30 ppm	75 g per 500 l water 150 g per 1000 l water	6 sachets per 500 l water 12 sachets per 1000 l water
50 g per 100 l water	100 ppm	250 g per 500 l water 500 g per 1000 l water	20 sachets per 500 l water 40 sachets per 1000 l water
5 g per 1000 l water	1 ppm	25 g per 5000 l water	2 sachets per 5000 l water

**APPLICATION RATES**

Crop / Use	Dosage Rate	Remarks
<b>Grapes</b>		<b>SuBana Grapes:</b> Apply in 1300 to 1800 litres water per hectare. Ensure thorough wetting of bunches and enclosing leaves. Care must be taken with wetting agents as russeting can occur. Both spray and dip methods can be applied. The correct timing of the application is critical for the success of the treatment.
<b>Sultanas (seedless)</b> Berry thinning (1)	5 g /100 l water	<b>First application:</b> When 50 % capsule has occurred in at least half of all bunches (half of all the bunches are 50 % in bloom).
Berry thinning (2)	5 g /100 l water	<b>Second application:</b> When 80 % capsule has occurred in at least half of all the bunches (half of all the bunches are 80 % in bloom).
<b>Lower Orange River area</b> To enlarge berries	15 g /100 l water plus non-ionic wetting agent	<b>First application:</b> Apply when berries have set and 50 % are ± 4 to 5 mm in diameter.
	15 g /100 l water plus non-ionic wetting agent	<b>Second application:</b> Apply when 75 % of berries are ± 4 to 5 mm in diameter.
	15 g /100 l water plus non-ionic wetting agent	<b>Third application:</b> Apply when 100 % of berries have are ± 4 to 5 mm in diameter.
<b>All other areas</b> To enlarge berries	15 g /100 l water plus non-ionic wetting agent	<b>First application:</b> Apply when berries have set and are 5 mm in diameter.
	15 g /100 l water plus non-ionic wetting agent	<b>Second application:</b> Apply 7 days after first application. (Application must not occur later than the post-berry stage.)
<b>Waltham Cross</b> Increase berry size and obtain a more even size berry	2,5 g /100 l water plus non-ionic wetting agent	Apply when Waltham Cross berries are 4 to 5 mm in diameter. Fill a container, which is large enough to hold one bunch with FALGRO® 20 SP solution and dip the entire bunch. Ensure that all berries are well wetted.
<b>Flame Seedless</b> Lower Orange River area Berry thinning	2,5 g /100 l water	Flame Seedless has naturally straggly bunches and therefore thinning may not be required in some instances. Apply during flowering when 80% in full bloom.
<b>Flame Seedless</b> Lower Orange River area To enlarge berries (1)	15 g /100 l water plus non-ionic wetting agent	<b>First Application:</b> Apply when berries are ± 7 to 8 mm in diameter.
To enlarge berries (2)	15 g /100 l water plus non-ionic wetting agent	<b>Second Application:</b> Apply when berries are ± 8 to 9 mm in diameter.
<b>Hops</b> To improve yield by enhancing growth in spring and for a shorter and more even hop treising period.	5 g /100 l water plus non-ionic wetting agent	Apply as a full cover spray during early spring; during September to middle October.
<b>Mangoes</b> To inhibit flowering of young trees.	50 g /100 l water plus non-ionic wetting agent	Before FALGRO® 20 SP is applied, break off all lower panicles. Apply as a full cover spray during the last week of July or during the first week of August.