



falgro[®]

PLANT GROWTH REGULATOR

Active Ingredient:

Contains 133g/kg gibberellic acid in the form of a tablet.

Each tablet contains 1g of gibberellins
comprising not less than 90% gibberellic acid
and not more than 10% other biologically active gibberellins.

Registered pursuant to the ACVM Act 1997 No.P5516
See www.foodsafety.govt.nz for registration conditions

Approved pursuant to the HSNO Act 1996 Approval code HSR000748
See www.epa.govt.nz for approval controls

Net Contents: 75g (10 tablets)

Distributed by:

Fruitfed Supplies

Building 2, 1 William Pickering Drive,
Albany, Auckland 0632

Tel: (09) 448 0510

Fax: (09) 448 0511

Part of PGG Wrightson Limited

For specialist advice in an emergency only
call, (0800 243 6225)

@Falgro is a registered trademark
of Fine Agrochemicals Ltd.

fine

Fine Agrochemicals Limited
Hill End House
Whitlington
Worcester
WR5 2RQ
UK

T +44 1905 361800

F +44 1905 361810

enquire@fine-agrochemicals.com

www.fine-agrochemicals.com

FAL/TAB/NZ/14



PLEASE READ THE LABEL COMPLETELY BEFORE USE

HSNO CLASSIFICATIONS: 9.1D, 9.2D

ECOTOXIC

HARMFUL TO AQUATIC ORGANISMS. Avoid contamination of any water supply with product or empty container.
HARMFUL TO THE SOIL ENVIRONMENT.

HANDLING PRECAUTIONS

EQUIPMENT: Apply using accurately calibrated equipment in accordance with The New Zealand Standard for the Management of Agrichemicals (NZS8409).

STORAGE: Keep out of reach of children. Store in original container, tightly closed in a locked, dry, cool location below 32°C. Use all solutions of Falgro on the day of preparation. Storage must be in accordance with current version of NZS 8409 Management of Agrichemicals. When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

PERSONAL PROTECTION: May cause eye irritation and mild skin irritation. Avoid contact with eyes and skin. When mixing or applying wear protective clothing including water proof gloves and goggles. Wash hands and exposed skin before meals and after work.

PRODUCT DISPOSAL: Dispose of this product only by using according to the label or at an approved landfill or other approved facility. Avoid contamination of any water supply with product.

CONTAINER DISPOSAL: Triple rinse container and add rinsate to spray tank. Burn in an appropriate incinerator, if circumstances such as wind direction permit. Otherwise crush or puncture and bury in a suitable landfill, or if appropriate recycle. Avoid contamination of any water supply with chemical or empty container.

FIRST AID

If swallowed do Not induce vomiting. For advice contact the National Poisons Center 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. Do not scrub the skin. If splashed in the eyes, wash out immediately with water

SPILLAGE: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely in approved landfill.

PRODUCT INFORMATION

Falgro contains gibberellic acid, (GA₃), a natural plant growth regulator derived by fermentation that occurs normally in many species of plant.

DIRECTIONS FOR USE

Mixing

After determining the rate of Falgro needed for the crop, add the correct number of tablets for the required tank volume and strength of spray, into approximately 2 litres of water to produce a concentrate solution. When the tablets have dissolved, add the solution, and the required rate of surfactant to the water in the spray tank, and agitate. For citrus the ideal spray solution pH is 4-5.5 to maximise GA₃ uptake.

Cherries:

To increase cherry firmness and quality apply approximately three weeks prior to the expected harvest date, when the cherries are straw coloured. Use at 10 ppm (10 tablets per 1000 litres) plus non-ionic surfactant at 250 ml per 1000 litres of water. Use high water rate to ensure thorough coverage of the crop, (1500-2000 litres/ha) in high volume sprays. Apply during cool conditions or in the evening to trees not under water stress.

Citrus:

Satsuma Mandarins: To reduce fruit numbers, apply in mid-late June at 25 ppm (25 tablets per 1000 litres) plus non-ionic surfactant at 250 ml/1000 litres of water. Use high water rates to ensure full coverage is obtained. (2000-2500 litres/ha). Do not apply if a heavy crop has been carried in the previous season, as flower numbers will be naturally low and the application of Falgro in this case, can stop flowering completely.

Navel Oranges: Rind defects: apply shortly after the end of natural fruit drop, (mid-February – late March) at 10 ppm. (10 tablets per 1000 litres) plus a non-ionic surfactant at 250 ml/1000 litres. Use high water rates to ensure full coverage is obtained. (2000-6000 litres/ha).

Other Uses: Falgro is known to be effective in many other applications. At rates from 10-100 ppm it had resulted in stem elongation, breaking of seed dormancy, earlier flowering, and set of fruit under adverse weather conditions when applied to a wide range of annual and perennial crops and ornamental plants, e.g. seedless grapes and zantedeschia lilies.

COMPATABILITY

Falgro should not be combined with oil sprays and should not be applied within four weeks of an oil spray as uptake will be negatively effected. Falgro is compatible with 2,4-D.

WITHHOLDING PERIOD: nil.

CONDITIONS OF SUPPLY - Read before buying and using.

If these conditions are unacceptable, return at once unopened. This product is of a high grade and conforms to the chemical description on the label. However, all warranties of any kind, expressed or implied, with respect to the fitness for purpose (including crop injury, ineffectiveness or any other unintended consequence), merchantable quality or any other matter, are expressly excluded. To the extent permitted by law, buyers and/or users accept that neither Fine Agrochemicals as the manufacturer, Fruited Supplies (part of PGG Wrightson Ltd) as distributor, nor any other of its distributors, accept any liability (including in negligence, tort and equity) or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use. The Terms of Sale that apply to this product are located at www.pggwrightson.co.nz > Our Company > Terms & Conditions > Terms of Sale.

® Falgro is a registered trademark of Fine Agrochemicals Ltd.

Distributed by:

Fruited Supplies

FRUITFED SUPPLIES
Building 2, 1 William Pickering Drive,
Albany, Auckland 0632.
T (09) 448 0510 F (09) 448 0511
Part of PGG Wrightsons Limited

fine

FINE AGROCHEMICALS LIMITED
Hill End House Whittington
Worcester WR5 2RQ UK
T +44 1905 361800 F +44 1905 361810
E enquire@fine-agrochemicals.com
www.fine-agrochemicals.com

For specialist advice in an emergency only, call (0800 243 6225)

FALNZ/15L