

**HSNO CLASSIFICATIONS:**  
No HSNO Classification triggered

**HANDLING PRECAUTIONS**

**EQUIPMENT:** Apply using accurately calibrated equipment in accordance with The New Zealand Standard for the Management of Agrichemicals (NZS 8409).

**STORAGE:** Keep out of reach of children. Store in original container, tightly closed in a locked, dry, cool well ventilated area. Storage should be in accordance with current version of NZS 8409 Management of Agrichemicals. When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

**PERSONAL PROTECTION:** When handling wear protective clothing, gloves and goggles. Wash hands and exposed skin before meals and after work.

**PRODUCT DISPOSAL:** Dispose of this product only by using according to the label or at an appropriate landfill. Avoid contamination of any water supply with product.

**CONTAINER DISPOSAL:** Triple rinse container and add rinsate to spray tank. Burn in an appropriate incinerator, if circumstances such as wind direction permit. Otherwise crush or puncture and bury in a suitable landfill, or if appropriate recycle. Avoid contamination of any water supply with chemical or empty container.

**FIRST AID**

If swallowed do Not induce vomiting. For advice contact the National Poisons Center 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. Do not scrub the skin. If splashed in the eyes, wash out immediately with water.

**SPILLAGE:** Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely in suitable landfill.

It is an offence to use this product on animals.

**DIRECTIONS FOR USE**

**Novagib** contains gibberellins A<sub>4</sub> and A<sub>7</sub>, natural plant growth regulators derived by fermentation, that occur normally in apples. When applied to developing fruit, **Novagib** improves the quality and marketability of the final harvested crop by significantly reducing the incidence of russetting and improving skin finish and appearance. This is achieved by increasing the growth rate and changing the morphology of the epidermis of the growing fruit.

**DATE OF MANUFACTURE:** As marked.

**BATCH NO.:** As marked.

©Novagib is a registered trademark of

**fine**

FINE AGROCHEMICALS LIMITED  
Hill End House Whittington  
Worcester WR5 2RQ UK

E enquire@fine.eu  
www.fine.eu



 **novagib**®

**A GROWTH REGULATOR FOR THE REDUCTION OF RUSSETING AND CRACKING IN APPLES**


**ADVANCED RUSSET CONTROL**

Active Ingredient: Contains 10g/litre of gibberellins A<sub>4</sub> and A<sub>7</sub> in the form of a soluble concentrate.

Registered pursuant to the ACVM Act, 1997 No. P5404.

See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

**NET CONTENTS: 1 Litre**



For specialist advice in an emergency only, call (0800 243 6225)



**RATES OF USE**

**Cox's Orange:** Apply 50 ml of **Novagib** per 100 litres of water at completion of bloom to petal fall followed by 2 or 3 further applications made at 10 day intervals.

**Other Varieties:** (e.g. Royal Gala, Pacific Rose, Fuji): Apply 50-100 ml of **Novagib** per 100 litres of water at completion of bloom to petal fall followed by 2 or 3 further applications made at 10 day intervals. Use the higher rate and greater number of applications where the risk of russetting is greater. Best effect will always be achieved by higher rates. Note: Rates higher than 10 ppm can result in a reduction of flowering the following season.

**WATER VOLUME**

Apply using an orchard sprayer in water volume sufficient to achieve complete coverage of fruitlets and penetration of target trees, usually 1000-1200 litres/ha. Spraying in still humid conditions will enhance the efficacy of **Novagib**.

**MIXING**

Add the required amount of **Novagib** to the half filled sprayer tank whilst agitating. Fill the tank and continue whilst spraying.

**COMPATIBILITY**

Consult your agrochemical advisor for advice on the use of **Novagib** in tank-mixture with other agrochemicals. It is usually best to apply **Novagib** alone.

**CONDITIONS OF SUPPLY - Read before buying and using.**

If these conditions are unacceptable, return at once unopened.

This product is of a high grade and conforms to the chemical description on the label. However, all warranties of any kind, expressed or implied, with respect to the fitness for purpose (including crop injury, ineffectiveness or any other unintended consequence), merchantable quality or any other matter, are expressly excluded. To the extent permitted by law, buyers and/or users accept that neither Fine Agrochemicals Ltd. as the manufacturer, Fruitfed Supplies (part of PGG Wrightson Ltd) as distributor, nor any other of its distributors, accept any liability (including in negligence, tort and equity) or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use. The Terms of Sale that apply to this product are located at [www.pggwrightson.co.nz](http://www.pggwrightson.co.nz) > Our Company> Terms & Conditions > Terms of Sale.

Distributed by:



Building 2  
1 William Pickering Drive  
Albany  
Auckland 0632  
Tel: (09) 448 0510, Fax: (09) 448 0511  
Part of PGG Wrightson Ltd.

NOV/NZ/1/L/14